

Maklumat tambahan indikasi untuk upload pada laman web

Year 2012

Products Approved For Additional Indication (DCA 250 – 29 MARCH 2012)

NO	PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1.	<p>1.1 Sandostatin LAR 10mg (Octreotide 10mg)</p> <p>1.2 Sandostatin LAR 20mg (Octreotide 20mg)</p> <p>1.3 Sandostatin LAR 30mg (Octreotide 30mg)</p>	<p>➤ INDICATION:</p> <ul style="list-style-type: none"> <i>Treatment of patients with advanced neuroendocrine tumours of the midgut; or of unknown primary origin where non-midgut sites of origin have been excluded.</i> <p>➤ POSOLOGY:</p> <p><i>Treatment of patients with advanced neuroendocrine tumours of the midgut; or of unknown primary origin where non-midgut sites of origin have been excluded.</i></p> <ul style="list-style-type: none"> <i>The recommended dose of Sandostatin LAR is 30mg administered every 4 weeks. Treatment with Sandostatin LAR for tumour control should be continued in the absence of tumour progression.</i> 	<p>NOVARTIS CORPORATION (MALAYSIA) SDN. BHD. LEVEL 15, CREST, 3 TWO SQUARE NO. 2, JALAN 19/1, 46300 PETALING JAYA, SELANGOR .</p>
2.	<p>Galvus 50mg Tablet (Vildagliptin 50mg)</p>	<p>➤ INDICATION:</p> <p><i>Galvus is indicated in the treatment of type 2 diabetes mellitus: As monotherapy</i></p> <ul style="list-style-type: none"> <i>in patients inadequately controlled by diet and exercise alone and for whom metformin is inappropriate due to contraindications or intolerance.</i> <p>➤ POSOLOGY:</p> <ul style="list-style-type: none"> <i>When used as monotherapy or in dual combination with metformin or a thiazolidinedione, the recommended daily dose of vildagliptin is 100mg, administered as one dose of 50 mg in the morning and one dose of 50 mg in the evening.</i> 	<p>NOVARTIS CORPORATION (MALAYSIA) SDN. BHD. LEVEL 15, CREST, 3 TWO SQUARE NO. 2, JALAN 19/1, 46300 PETALING JAYA, SELANGOR .</p>